

**For Research Use Only.
Not for use in diagnostic procedures.**



Anti-V5-tag mAb-HRP-Direct

CODE No.	M215-7
CLONALITY	Monoclonal
CLONE	OZA3
ISOTYPE	Mouse IgG2b κ
QUANTITY	100 μ L
SOURCE	Purified IgG from hybridoma supernatant
IMMUNOGEN	Carrier protein conjugated synthetic peptide, GKPIPPLLGLDST (V5-tag)
FORMURATION	PBS/Preservative/Stabilizer
STORAGE	This antibody solution is stable for one year from the date of purchase when stored at 4°C.
APPLICATION-CONFIRMED	
<u>Western blotting</u>	1:1,000 for chemiluminescence detection system

For more information, please visit our web site <http://ruo.mbl.co.jp/>



RELATED PRODUCTS

Antibodies

M048-3	Anti-GFP mAb (1E4)
D153-3	Anti-GFP mAb (RQ2)
D153-6	Anti-GFP mAb-Biotin (RQ2)
D153-8	Anti-GFP mAb-Agarose (RQ2)
598	Anti-GFP pAb (polyclonal)
598-7	Anti-GFP pAb-HRP-Direct (polyclonal)
PM073	Anti-Renilla GFP pAb (polyclonal)
M208-3	Anti-RFP mAb Cocktail (1G9, 3G5)
M155-3	Anti-RFP mAb (8D6)
M165-3	Anti-RFP mAb (3G5)
M165-8	Anti-RFP mAb-Agarose (3G5)
M204-3	Anti-RFP mAb (1G9)
M204-7	Anti-RFP mAb-HRP-Direct (1G9)
PM005	Anti-RFP pAb (polyclonal)
PM005-7	Anti-RFP pAb-HRP-Direct (polyclonal)
M180-3	Anti-HA-tag mAb (TANA2) (200 µL)
M180-6	Anti-HA-tag mAb-Biotin (TANA2)
M180-7	Anti-HA-tag mAb-HRP-Direct (TANA2)
561	Anti-HA-tag pAb (polyclonal) (0.1 mL)
561-7	Anti-HA-tag pAb-HRP-Direct (polyclonal)
561-8	Anti-HA-tag pAb-Agarose (polyclonal)
M132-3	Anti-HA-tag mAb (5D8)
M185-3L	Anti-DDDDK-tag mAb (FLA-1) (1 mL)
M185-7	Anti-DDDDK-tag mAb-HRP-Direct (FLA-1)
PM020	Anti-DDDDK-tag pAb (polyclonal)
PM020-7	Anti-DDDDK-tag pAb-HRP-Direct (polyclonal)
PM020-8	Anti-DDDDK-tag pAb-Agarose (polyclonal)
M192-3	Anti-Myc-tag mAb (My3) (200 µL)
M192-6	Anti-Myc-tag mAb-Biotin (My3)
M047-3	Anti-Myc-tag mAb (PL14)
M047-6	Anti-Myc-tag mAb-Biotin (PL14)
M047-7	Anti-Myc-tag mAb-HRP-Direct (PL14)
M047-8	Anti-Myc-tag mAb-Agarose (PL14)
562	Anti-Myc-tag pAb (polyclonal) (0.1 mL)
D291-3	Anti-His-tag mAb (OGHis) (200 µL)
D291-6	Anti-His-tag mAb-Biotin (OGHis)
D291-7	Anti-His-tag mAb-HRP-Direct (OGHis)
D291-8	Anti-His-tag mAb-Agarose (OGHis)
D291-A48	Anti-His-tag mAb-Alexa Fluor [®] 488 (OGHis)
D291-A59	Anti-His-tag mAb-Alexa Fluor [®] 594 (OGHis)
D291-A64	Anti-His-tag mAb-Alexa Fluor [®] 647 (OGHis)
M089-3	Anti-His-tag mAb (6C4)
M136-3	Anti-His-tag mAb (2D8)
PM032	Anti-His-tag pAb (polyclonal)
PM032-8	Anti-His-tag pAb-Agarose (polyclonal)
M167-3	Anti-V5-tag mAb (1H6)
M215-3	Anti-V5-tag mAb (OZA3)
M215-7	Anti-V5-tag mAb-HRP-Direct (OZA3)
PM003	Anti-V5-tag pAb (polyclonal)
PM003-7	Anti-V5-tag pAb-HRP-Direct (polyclonal)
PM003-8	Anti-V5-tag pAb-Agarose (polyclonal)
PM021	Anti-S-tag pAb (polyclonal)
PM070	Anti-E-tag pAb (polyclonal)
PM022	Anti-T7-tag pAb (polyclonal)
563	Anti-VSV-G-tag pAb (polyclonal)
M071-3	Anti-GST-tag mAb (3B2)
M209-3	Anti-GST-tag mAb (GT5)
PM022	Anti-GST-tag pAb (polyclonal)

M095-3	Anti-Luciferase mAb (2D4)
PM016	Anti-Luciferase pAb (polyclonal)
PM047	Anti-Renilla Luciferase pAb (polyclonal)
M094-3	Anti-β-galactosidase mAb (5A3)
PM049	Anti-β-galactosidase pAb (polyclonal)
M091-3	Anti-MBP (Maltose Binding Protein) mAb (1G12)
M013-3	Anti-Thioredoxin (Trx-tag) mAb (2C9)
PM015	Anti-CBD (Chitin Binding Domain) pAb (polyclonal)
PM071	Anti-Calmodulin Binding Protein-tag pAb (polyclonal)
M211-3	Anti-Strep-tag II mAb (4F1)
M214-3	Anti-mini-AID-tag mAb (1E4)

Smart-IP series

3190	Magnetic Rack
M180-9	Anti-HA-tag mAb-Magnetic beads (TANA2)
M132-9	Anti-HA-tag mAb-Magnetic beads (5D8)
M185-9	Anti-DDDDK-tag mAb-Magnetic beads (FLA-1)
M047-9	Anti-Myc-tag mAb-Magnetic beads (PL14)
D291-9	Anti-His-tag mAb-Magnetic beads (OGHis)
D153-9	Anti-GFP mAb-Magnetic beads (RQ2)
M165-9	Anti-RFP mAb-Magnetic beads (3G5)
M198-9	Anti-E-tag mAb-Magnetic beads (21D11)
M167-11	Anti-V5-tag mAb-Magnetic Beads (1H6)
D058-9	Anti-Multi Ubiquitin mAb-Magnetic beads (FK2)
M075-9	Mouse IgG1 (isotype control)-Magnetic beads
M076-9	Mouse IgG2a (isotype control)-Magnetic beads
M077-9	Mouse IgG2b (isotype control)-Magnetic beads
M081-9	Rat IgG2a (isotype control)-Magnetic beads
M180-10	Anti-HA-tag mAb-Magnetic Agarose (TANA2)
M132-10	Anti-HA-tag mAb-Magnetic Agarose (5D8)
M185-10	Anti-DDDDK-tag mAb-Magnetic Agarose (FLA-1)
M047-10	Anti-Myc-tag mAb-Magnetic Agarose (PL14)
D291-10	Anti-His-tag mAb-Magnetic Agarose (OGHis)
D153-10	Anti-GFP mAb-Magnetic Agarose (RQ2)
M165-10	Anti-RFP mAb-Magnetic Agarose (3G5)
M167-10	Anti-V5-tag mAb-Magnetic Agarose (1H6)
M198-10	Anti-E-tag mAb-Magnetic Agarose (21D11)


Protein Purification Kits

3320	HA-tagged Protein PURIFICATION KIT
3321	HA-tagged Protein PURIFICATION GEL (1 mL)
3325	DDDDK-tagged Protein PURIFICATION KIT
3326	DDDDK-tagged Protein PURIFICATION GEL (1 mL gel, 5 mg peptide)
3328	DDDDK-tagged Protein PURIFICATION GEL (5 mL gel)
3325-205	DDDDK-tag peptide (1 mg x 5)
3326K	DDDDK-tagged Protein PURIFICATION CARTRIDGE (1 mL x 1)
3305	c-Myc-tagged Protein MILD PURIFICATION KIT
3306	c-Myc-tagged Protein MILD PURIFICATION GEL (1 mL gel, 1 mg peptide)
3310	His-tagged Protein PURIFICATION KIT
3311	His-tagged Protein PURIFICATION GEL (1 mL gel, 5 mg peptide)
3315	V5-tagged Protein PURIFICATION KIT
3316	V5-tagged Protein PURIFICATION GEL (1 mL)

Other related antibodies and kits are also available.
Please visit our website at <http://ruo.mbl.co.jp/>

SDS-PAGE & Western blotting

- 1) Mix the sample with equal volume of Laemmili's sample buffer.
- 2) Boil the sample for 2-3 min. and centrifuge. Load the sample per lane in a 1-mm-thick SDS-polyacrylamide gel (12.5% acrylamide) for electrophoresis.
- 3) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at 1 mA/cm² for 1 hr. in a semi-dry transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% MeOH). See the manufacturer's manual for precise transfer procedure.
- 4) To reduce nonspecific binding, soak the membrane in 10% skimmed milk (in PBS, pH 7.2) over night at 4°C.
- 5) Wash the membrane with PBS-T [0.05% Tween-20 in PBS] (5 min. x 3 times).
- 6) Incubate the membrane with primary antibody diluted with 1% skimmed milk (in PBS, pH 7.2) as suggested in the **APPLICATIONS** for 1 hr. at room temperature. (The concentration of antibody will depend on the conditions.)
- 7) Wash the membrane with PBS-T (5 min. x 3 times).
- 8) Wipe excess buffer on the membrane, and then incubate it with appropriate chemiluminescence reagent for 1 min. Remove extra reagent from the membrane by dabbing with paper towel, and seal it in plastic wrap.
- 9) Expose to an X-ray film in a dark room for 1 min. Develop the film as usual. The condition for exposure and development may vary.




Western blot analysis of V5-tagged proteins

Lane 1: V5-tagged TPO in insect cell culture sup (20 μ L/lane)

Lane 2: V5-tagged GFP (25 ng/lane)

Lane 3: V5-tagged β -galactosidase/293T (10 μ L/lane)

Immunoblotted with Anti-V5-tag mAb-HRP-Direct (M215-7)



Western blot analysis of V5-tagged protein

Sample: Insect cell culture sup containing V5-tagged TPO

Lane 1: IP with Mouse IgG2b (M077-3)

Lane 2: IP with Anti-V5-tag mAb (M215-3)

Immunoblotted with Anti-V5-tag mAb-HRP-Direct (M215-7)