

# Anti-CD81 (TAPA1) mAb-Biotin

<b>CODE No.</b>	MEX003-6
<b>CLONALITY</b>	Monoclonal
<b>CLONE</b>	A103-10
<b>ISOTYPE</b>	Mouse IgG2a κ
<b>QUANTITY</b>	50 µL, 1 mg/mL
<b>SOURCE</b>	Purified IgG from hybridoma supernatant
<b>IMMUNOGEN</b>	Human prostate carcinoma cell line (PC3) derived exosomes (prepared by ultracentrifugation from cultured supernatant)
<b>FORMULATION</b>	PBS containing 1% BSA and 0.1% ProClin 950
<b>STORAGE</b>	This antibody solution is stable for one year from the date of purchase when stored at 4°C.

## APPLICATIONS-CONFIRMED

<u>Western blotting</u>	1 µg/mL for chemiluminescence detection system ( <u>non-reducing condition</u> )
<u>Flow cytometry</u>	5-10 µg/mL
<u>Sandwich CLEIA</u>	Can be used. Please refer to " <u>ExoDiluent for Immunoassay (MEX1001)</u> " together.
<u>Exosome isolation</u>	Can be used. Please refer to " <u>ExoCap™ Streptavidin Kit (MEX-SA)</u> " together.

## APPLICATION-UNDER EVALUATION

<u>Sandwich ELISA</u>	Can be used.
-----------------------	--------------

## SPECIES CROSS REACTIVITY on WB

Species	Human	Monkey	Mouse	Rat	Hamster
Cells	HEK293T	Not tested	WR19L	Not tested	Not tested
Reactivity	+		-		

<b>Entrez Gene ID</b>	975 (Human)
-----------------------	-------------

## REFERENCES

- 1) Melo, S. A., *et al.*, *Nature* **523**, 177-182 (2015)
- 2) Yoshioka, Y., *et al.*, *Nat. Commun.* **5**, 3591 (2014)
- 3) Pols, M. S. and Klumperman, J., *Exp. Cell Res.* **315**, 1584-1592 (2009)
- 4) Simons, M. and Raposo, G., *Curr. Opin. Cell Biol.* **21**, 575-581 (2009)

For more information, please visit our web site <http://ruo.mbl.co.jp/>



MEDICAL & BIOLOGICAL LABORATORIES CO., LTD.

URL <http://ruo.mbl.co.jp/>

e-mail [support@mbl.co.jp](mailto:support@mbl.co.jp), TEL 052-238-1904

## RELATED PRODUCTS

### Antibodies

MEX001-3	Anti-CD9 mAb (A100-4)
MEX001-4	Anti-CD9 mAb-FITC (A100-4)
MEX001-6	Anti-CD9 mAb-Biotin (A100-4)
MEX001-12	Anti-CD9 mAb-ALP (A100-4)
MEX002-3	Anti-CD63 (LAMP-3) mAb (C047-1)
MEX002-4	Anti-CD63 (LAMP-3) mAb-FITC (C047-1)
MEX002-6	Anti-CD63 (LAMP-3) mAb-Biotin (C047-1)
MEX002-12	Anti-CD63 (LAMP-3) mAb-ALP (C047-1)
MEX003-3	Anti-CD81 (TAPA1) mAb (A103-10)
MEX003-4	Anti-CD81 (TAPA1) mAb-FITC (A103-10)
MEX003-6	Anti-CD81 (TAPA1) mAb-Biotin (A103-10)
MEX003-12	Anti-CD81 (TAPA1) mAb-ALP (A103-10)
MEX004-6	Anti-EpCAM mAb-Biotin (B8-4)
K0142-6	Anti-PSMA (Human) mAb-Biotin (107-1A4)
D281-6	Anti-CD61 (GPIIIa) (Human) mAb-Biotin (T74)
M232-6	Anti-CD36 (GPIV) (Human) mAb-Biotin (1-3)
M232-3	Anti-CD36 (GPIV) (Human) mAb (1-3)
D252-3	Anti-CD9 (Human) mAb (10H6)
D252-5	Anti-CD9 (Human) mAb-PE (10H6)
D131-3	Anti-CD9 (Mouse) mAb (JF9)
D131-4	Anti-CD9 (Mouse) mAb-FITC (JF9)
D263-3	Anti-CD63 (LAMP-3) (Mouse) mAb (R5G2.1)
D082-3	Anti-CD151 (SFA-1) (Human) mAb (SFA1.2B4)
D082-5	Anti-CD151 (SFA-1) (Human) mAb-PE (SFA1.2B4)
D050-3	Anti-CD29 (Integrin $\beta$ 1) (Human) mAb (AG89)
D050-5	Anti-CD29 (Integrin $\beta$ 1) (Human) mAb-PE (AG89)
D276-3	Anti-CD36 (GPIV) (Human) mAb (GS95)
D276-A48	Anti-CD36 (GPIV) (Human) mAb -Alexa Fluor <sup>®</sup> 488 (GS95)
D276-A64	Anti-CD36 (GPIV) (Human) mAb -Alexa Fluor <sup>®</sup> 647 (GS95)
D269-3	Anti-EpCAM (CD326) (Mouse) mAb (2-17-F1)
M076-6	Mouse IgG2a (isotype control)-Biotin (6H3)
M076-12	Mouse IgG2a (isotype control)-ALP (6H3)

### Kits

MEX-SA	ExoCap™ Streptavidin Kit
MEX-SA123	ExoCap™ Streptavidin CD9/CD63/CD81 Set
MEX-E	ExoCap™ Nucleic Acid Elution Buffer
MEX1001	ExoDiluent for Immunoassay
RN1001	RIP-Assay Kit

### Others (reagents, equipments, etc.)

MJS002V2	Protein G-Magnetic Beads
3190	Magnetic Rack


Other related antibodies and kits are also available.

Please visit our website at <http://ruo.mbl.co.jp/>

### **SDS-PAGE & Western blotting (non-reducing condition)**

- 1) Wash  $1 \times 10^7$  cells 3 times with PBS and suspend with 1 mL of Laemmli's sample buffer (non-reducing condition), then sonicate briefly (up to 10 sec.)
- 2) Boil the samples for 3 min. and centrifuge. Load 10  $\mu\text{L}$  of the sample per lane in a 1-mm-thick SDS-polyacrylamide gel (15% acrylamide) for electrophoresis.
- 3) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at 10 V for 50 min. in a semi-dry transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% MeOH). See the manufacturer's manual for precise transfer procedure.
- 4) To reduce nonspecific binding, soak the membrane in 5% skimmed milk (in PBS, pH 7.2) overnight at 4°C.
- 5) Wash the membrane with PBS-T [0.05% Tween-20 in PBS] (5 min.  $\times$  3 times).
- 6) Incubate the membrane with primary antibody diluted with 1% skimmed milk (in PBS, pH 7.2) as suggested in the **APPLICATIONS** for 1 hr. at room temperature. (The concentration of antibody will depend on the conditions.)
- 7) Wash the membrane with PBS-T (5 min.  $\times$  3 times).
- 8) Incubate the membrane with the 1:20,000 Streptavidin-Horseradish Peroxidase (GE Healthcare; code no. RPN4401) diluted with 1% skimmed milk (in PBS, pH 7.2) for 1 hr. at room temperature.
- 9) Wash the membrane with PBS-T (5 min.  $\times$  3 times)
- 10) Wipe excess buffer on the membrane, then incubate it with appropriate chemiluminescence reagent for 1 min. Remove extra reagent from the membrane by dabbing with paper towel, and seal it in plastic wrap.
- 11) Expose to an X-ray film in a dark room for 10 min. Develop the film as usual. The condition for exposure and development may vary.

(Positive control for Western blotting; HEK293T)



### ***Western blot analysis of CD81 (TAPA1) protein***


Lane 1: HEK293T  
Lane 2: WR19L

Immunoblotted with Anti-CD81 (TAPA1) mAb-Biotin (MEX003-6)

### **Flow cytometric analysis**

- 1) Wash the cells ( $5 \times 10^5$  cells/sample) 1 time with 1 mL of washing buffer [PBS containing 2% fetal calf serum (FCS)].
- 2) Add 10  $\mu$ L of Clear Back (human Fc receptor blocking reagent, MBL; code no. MTG-001) to the cell pellet after tapping. Mix well and incubate for 10 min. at room temperature.
- 3) Add 50  $\mu$ L of the primary antibody at the concentration as suggested in the **APPLICATION** diluted with washing buffer.
- 4) Mix well and incubate for 20 min. at room temperature.
- 5) Wash the cells 1 time with 1 mL of washing buffer.
- 6) Add FITC-conjugated streptavidin diluted with washing buffer. Mix well and incubate for 20 min. at room temperature.
- 7) Wash the cells 1 time with 1 mL of washing buffer.
- 8) Resuspend the cells with 500  $\mu$ L of the washing buffer and analyze by a flow cytometer.

(Positive controls for Flow cytometry; HeLa and HEK293T)




### ***Flow cytometric detection of human CD81 (TAPA1)***

Left: HeLa  
Right: HEK293T

Open: Anti-CD81 (TAPA1) mAb-Biotin (MEX003-6)  
Closed: Mouse IgG2a (isotype control)-Biotin (M076-6)

**Sandwich CLEIA (chemiluminescence enzyme immunoassay)**




**Sandwich CLEIA for measurement of CD81 (TAPA1) expressed HeLa-derived exosomes**

Sample: HeLa-derived exosomes prepared by ultracentrifugation

Capture Antibody: Anti-CD9 mAb (MEX001-3)

Detection Antibody: Anti-CD81 (TAPA1) mAb-Biotin (MEX003-6)



**Sandwich CLEIA for measurement of CD81 (TAPA1) expressed exosomes from human serum**

Sample: Pooled human serum from healthy volunteers

Capture Antibody: Anti-CD9 mAb (MEX001-3)

Detection Antibody: Anti-CD81 (TAPA1) mAb-Biotin (MEX003-6)